

We Claim:

- Sub B1 D1
1. A method for microdissecting tissue, comprising:
labeling a sample of the tissue by exposing the tissue to a sufficient
concentration of a fluorescent specific binding agent for a sufficiently short period of
time to reduce a binding time of the agent to the tissue to reduce loss of a biological
molecule in the tissue; and
identifying a component of interest in the sample to which the fluorescent
specific binding agent binds by detecting fluorescence of the component of interest in
the tissue; and
microdissecting components of interest from the tissue.
2. The method of claim 1, wherein identifying the component comprises
intensifying a signal from the fluorescent specific binding agent to provide an
intensified image signal.
3. The method of claim 2, wherein microdissecting comprises dissecting tissue
components with a laser beam, and the method further comprises selectively filtering an
image of the laser beam to reduce laser-induced distortion of the intensified image.
4. The method of claim 1, wherein the specific binding agent comprises an
aqueous solution, and the biomolecule is RNA, DNA or a protein, which is lost in the
presence of water.
5. The method of claim 4, wherein the biomolecule is RNA.
6. The method of claim 1, wherein microdissecting comprises applying a
capture member to the sample of tissue, and applying laser energy to the component of
interest to adhere the component to the capture member.
- Sub B2
- RECEIVED

Sub B3 Di
7. The method of claim 1, wherein the sufficient concentration of fluorescent specific binding agent is sufficient to avoid loss of more than about 5% of the biomolecule.

5 8. The method of claim 7, wherein the sufficient concentration of fluorescent specific binding agent is sufficient to avoid loss of more than about 10% of the biomolecule.

Sub B4 Di
9. The method of claim 1, wherein the fluorescent specific binding agent includes a fluorescent antibody, lectin, protein A, protein G and mixtures thereof.

10 10. The method of claim 1, wherein the sufficient concentration of specific binding agent is at least 0.02 mg/mL

15 11. The method of claim 10 wherein the sufficient concentration of specific binding agent is at least 0.1 mg/mL.

12. The method of claim 9, further comprising pre-mixing a primary antibody and a secondary antibody, at least one of which is fluorescent, prior to exposing the tissue to the specific binding agent to reduce a time of exposure of the tissue to the specific binding agent.

13. The method of claim 2, wherein the fluorescent specific binding agent is present in a sufficient concentration that, when the tissue is exposed to the fluorescent specific binding agent for less than about 5 minutes, the intensified image signal is detectable.

Sub B5 Di
14. The method of claim 13, wherein the fluorescent specific binding agent is present in a sufficient concentration that, when the tissue is exposed to the fluorescent

specific binding agent for less than about 3 minutes, the intensified image signal is detectable.

15. The method of claim 14, wherein the fluorescent specific binding agent is present in a sufficient concentration that, when the tissue is exposed to the fluorescent specific binding agent for not more than about 1 minute, the intensified image signal is detectable.

16. The method of claim 3, wherein microdissecting comprises targeting tissue components with a target laser beam, and viewing the intensified image through an infrared filter that selectively minimizes image distortion caused by the laser beam, without eliminating the signal image.

17. A method of performing tissue microdissection of a tissue specimen, comprising:
exposing the tissue specimen to at least one fluorescently labeled antibody which specifically binds to a component of interest in the tissue, wherein the tissue is exposed to a sufficient concentration of the antibody, in an aqueous solution, for a sufficient period of time to label the component of interest without substantially degrading RNA in the tissue;

washing unbound antibody from the tissue;
intensifying an image of the tissue specimen which has been exposed to the fluorescently labeled antibody to obtain an intensified fluorescent signal from the tissue;

applying a transfer member to the tissue;
directing a target laser beam to the component of interest in the tissue, to mark the component that is to be dissected from the tissue specimen, while viewing the target laser beam through an infrared filter that selectively filters infrared radiation but not the fluorescent signal, to minimize heat distortion of the intensified image, while still viewing the intensified signal; and

C1
applying radiant laser energy to the component of interest to transfer the component to the transfer member.

18. The method of claim 17, wherein exposing the tissue to a sufficient concentration of the at least one fluorescently labeled antibody comprises exposing the tissue to a concentration of at least 0.04 mg/mL of the fluorescently labeled antibody.

19. The method of claim 18, wherein exposing the tissue to a sufficient concentration of the at least one fluorescently labeled antibody comprises exposing the tissue to a concentration of at least 0.10 mg/mL of the at least one fluorescently labeled antibody

20. The method of claim 17, wherein exposing the tissue to the at least one fluorescently labeled antibody comprises exposing the tissue to the at least one fluorescently labeled antibody for less than about 5 minutes.

21. The method of claim 20, wherein exposing the tissue to the at least one fluorescently labeled antibody comprises exposing the tissue to the at least one fluorescently labeled antibody for less than about 3 minutes.

22. The method of claim 21, wherein exposing the tissue to the fluorescently labeled antibody comprises exposing the tissue to the antibody for no more than about 1 minute.

23. An apparatus for rapid immunofluorescence laser capture microdissection, comprising:
a laser capture microdissection microscope;
a light source capable of exciting fluorescence from fluorescent specific binding agents disposed to illuminate a sample placed on the sample stage of the microscope;

an image intensifier disposed between the sample stage of the microscope and
an image acquisition system; and
an infrared filter disposed between the sample stage and the image intensifier.

24. A method for fluorescently staining a tissue section for microdissection, comprising:

fixing a tissue section with a non-crosslinking fixative;

10 rinsing the tissue section twice with an aqueous buffered solution for about 5 seconds per rinse;

incubating the fixed tissue section with an aqueous fluorescent specific binding agent solution of sufficient concentration to selectively label cells within the tissue section in about 1 minute.

15 rinsing the tissue section twice with an aqueous buffered solution for about 5 seconds per rinse;

dehydrating the tissue section; and

drying the tissue section

25. The method of claim 24, wherein the aqueous buffered solution is
20 diethylpyrocatechol-treated phosphate-buffered saline solution.

26. The method of claim 24 wherein the fluorescent specific binding agent solution comprises a primary antibody covalently linked to a fluorescent molecule.

25 27. The method of claim 24, wherein the fluorescent specific binding agent solution comprises a pre-mixed solution of primary antibody and fluorescently labeled secondary antibody.

28. The method of claim 24, wherein the fluorescent specific binding agent solution comprises a premixed solution of a fluorescently labeled primary antibody and a fluorescently labeled secondary antibody.

5 29. The method of claim 24, wherein the fluorescent specific binding agent solution comprises a solution of fluorescently labeled lectin.

10 30. The method of claim 24, wherein the fluorescent specific binding agent solution comprises a mixture of primary antibody and fluorescently labeled protein A or G.

31. The method of claim 30, wherein primary antibody is also fluorescently labeled.

15 32. The method of claim 24, wherein the non-crosslinking fixative is selected from the group consisting of ethanol, acetone, methanol and mixtures thereof.

20 33. The method of claim 24, wherein the fluorescent specific binding agent solution further comprises an enzyme inhibitor.

34. The method of claim 33, wherein the enzyme inhibitor is chosen from the group consisting of RNase inhibitor, DNase inhibitors, protease inhibitors and mixtures thereof.

25 ³⁵ 34. A method for immunofluorescently labeling tissue that preserves biological molecules, comprising:

contacting the tissue with aqueous fluorescent antibody solutions of sufficient concentration to selectively label immunophenotypically similar cells against which the fluorescent antibodies are directed in less than about five minutes.

30

Sub
B9
D1

WDN/TMH:jl 6/13/2001
Atty Ref. No. 4239-58570

E-133-00/0

Express Mail No. EL828141226US
Date of Deposit: June 13, 2001

35. The method of claim 34, wherein the aqueous fluorescent antibody solutions are of sufficient concentration to selectively label immunophenotypically similar cells against which the fluorescent antibodies are directed in less than about three minutes.

36. The method of claim 35, wherein the aqueous fluorescent antibody solutions are of sufficient concentration to selectively label immunophenotypically similar cells against which the fluorescent antibodies are directed in less than about three minutes.

Year	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100
1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	

Add DI